

EXCITATION AND DESTRUCTION OF THE OH⁺ and OD⁺ IONS by H ATOMS

Niyazi Bulut¹, François Lique² and Octavio Roncero³

¹University of Firat, Elazig, Turkey

²LOMC - UMR 6294, CNRS-Universit_e du Havre, 25 rue Philippe Lebon, BP 1123 - 76 063 Le Havre cedex, France

³Instituto de Fisica Fundamental (IFF-CSIC), C.S.I.C., Serrano 121-123, 28006 Madrid, Spain.

In space, deuterated species are surprisingly highly abundant. Following the recent discovery of OH⁺, we are studying the possible formation of the OD⁺ molecule in the interstellar medium. New quantum reactive scattering calculations for the O⁺ + HD and OD⁺ + H collisions have been carried out to obtain state-to-state cross sections and rate constants by means of an accurate wave packet approach [1-3] using the doublet and quartet ground H₂O⁺ electronic potential energy surfaces correlating to the open shell reactants. Calculations were performed for collision energies in the range of 1 meV to 0.7 eV and for different initial rotational excitation of the reagent molecules. These calculations provide us detailed information about the possible detection of OD⁺ in space. A review will be presented for ion-molecule collision processes that occur in the evolving chemistry of astronomical environments.

References:

- [1] N. Bulut, J. F. Castillo, P.G. Jambrina, J. Klos, O. Roncero, F. J. Aoiz, and L. Banares, J. Phys. Chem. A, 2015, 119, 11951-11962.
- [2] N. Bulut, F. Lique and O. Roncero, J. Phys. Chem. A, 2015, 119, 12082-12089.
- [3] F. Lique, N. Bulut, O. Roncero, MNRAS, 2016, 461, 4477-4481.